


PREMATURITY PROFILE WEST VIRGINIA


IN AN AVERAGE WEEK IN WEST VIRGINIA



PRETERM BIRTH RATE IN WEST VIRGINIA

PERCENTAGE OF LIVE BIRTHS

PRETERM BIRTH RATE, 2011-2020



- In 2020, there were 2,082 preterm births in West Virginia, representing 12.0% of live births.
- In 2020, 1 in 8 babies (12.0% of live births) was born preterm in West Virginia.

PRETERM RATES BY COUNTY, 2017-2020 AVERAGE



- Higher than WV Rate of 12.1 (25)
- **Between HP 2030 Objective and WV Rate (22)**
- Met or lower than HP 2030 Objective of 9.4 (8)

Every baby in West Virginia deserves the chance to be born healthy.

- In 2017-2020, 25 West Virginia counties had a preterm birth rate higher than the state rate of 12.1%.
- In 2017-2020, 8 West Virginia counties had met the March of Dimes preterm rate goal of 9.4%.



PREMATURITY PROFILE WEST VIRGINIA

PRETERM BIRTH BY RACE AND ETHNICITY

The March of Dimes is concerned about inequities in health and health care that contribute to higher rates of preterm birth among different racial and ethnic groups.

Hispanic 10.2 12.0 White 16.0 Black 10.3 Asian/ Pacific Islander 12.1 Total 0 2 4 6 8 10 12 14 16 18

PERCENTAGE OF PRETERM LIVE BIRTHS (2018-2020 AVERAGE)


- During 2018-2020 (average) in West Virginia, preterm birth rates were highest for black infants (16.0%), followed by Whites (12.0%), Asian/Pacific Islanders (10.3%) and Hispanics (10.2%).
- Black infants (16.0%) were about 2 times as likely as Hispanic infants (10.2%) to be born preterm during 2018-2020 (average).
 - Of all live births in West Virginia during 2018-2020 (average), 2.1% were Hispanic, 91.7% were white, 4.0% were black, 0.1% were American Indian/Alaska Native and 1.0% were Asian/Pacific Islander.

CONSEQUENCES OF PRETERM BIRTH

Preterm birth, along with low birth weight babies, make up the second leading cause of infant deaths after birth defects. Health consequences of preterm birth include developmental delays, chronic respiratory problems and vision and hearing impairment¹. Having a preterm baby impacts families emotionally and financially.



The annual societal economic cost (medical, educational, and lost productivity)³ associated with preterm birth in West Virginia is an estimated \$52,000.





RISK FACTORS FOR PRETERM BIRTH

Rather than having one cause, preterm birth seems to be triggered by multiple, interacting biologic and environmental factors. We do know that a history of preterm birth, multiple gestations, and certain uterine and/or cervical problems are the strongest risk factors for preterm birth. The factors below also contribute to preterm birth.⁴

MATERNAL AGE

During 2018-2020 (average) in West Virginia, preterm birth rates were highest for women ages 40 and older (16.9%), followed by women ages 30-39 (13.1%), ages 20-29 (11.6%) and under age 20 (11.4%).

BIRTH SPACING

Birth spacing, or inter-pregnancy interval, is the timing between a live birth and the beginning of the next pregnancy. Birth spacing of less than 18 months increases the risk of preterm birth and other adverse outcomes. In West Virginia, 32.9% of pregnancies with a prior live birth have a birth spacing of less than 18 months (2018-2020 average).

SMOKING

In 2020, 25.0% of women of childbearing age reported smoking in West Virginia. Smoking is an important determinant of health and a significant factor contributing to preterm births.

OBESITY

While obesity does not directly cause preterm birth, it does increase rates of medical complications (e.g., hypertension, diabetes) that contribute to preterm birth. In West Virginia, 37.6% of women of childbearing age were obese in 2020.

MULTIPLE BIRTHS

In West Virginia in 2020, 10.4% of singleton births were preterm, compared to 63.1% of multiple births. Multiple births represent 3.0% of live births in West Virginia. Current multifetal pregnancy is one of the most consistently identified risk factors for preterm birth.

HEALTH INSURANCE COVERAGE

In 2020, about 1 in 10 women of childbearing age (9.7%) was uninsured in West Virginia. Health care before, during and after pregnancy can help identify and manage conditions that contribute to preterm birth.

OTHER CONTRIBUTING FACTORS⁴

Other factors contributing to preterm birth include: infection (especially genito-urinary), diabetes mellitus, hypertension, late or no prenatal care, alcohol and illicit drug use, and social determinants of health. Socioeconomic status at both the individual and community level (e.g., income/poverty, job status, education) as well as psychosocial factors (e.g., chronic stress, lack of social support) are associated with an increased risk of preterm birth.



PREMATURITY PROFILE WEST VIRGINIA

MARCH OF DIMES MISSION

March of Dimes leads the fight for the health of all moms and babies. We support research, lead programs and provide education and advocacy so that every baby can have the best possible start. Building on a successful 80-year legacy of impact and innovation, we empower every mom and every family. Visit <u>https://www.marchofdimes.org/</u> or <u>https://nacersano.marchofdimes.org/</u> for more information.

FOOTNOTES

- 1. Preterm is less than 37 weeks of pregnancy.
- 2. Late preterm is between 34 and 36 weeks gestation.
- 3. Very preterm is less than 32 weeks.
- 4. All race categories exclude Hispanics.
- Smoking is defined as having ever smoked 100 cigarettes in a lifetime and currently smoking everyday or some days. Percent reported is among women ages 18-44.
- 6. Multiple deliveries include twin, triplet and higher order deliveries.
- Birth spacing is the period of time between giving birth and getting pregnant again. Its also called pregnancy spacing or interpregnancy interval (also called IPI).
- Obesity is defined as a Body Mass Index of 30 or more. Body Mass Index (BMI) is a number calculated from a person's weight and height. Percent reported is among women ages 18 -44.
- 9. A woman was considered uninsured if she was not covered by any type of health insurance at the time of the survey. Percent reported is among women ages 15-44.
- 10. Social determinants of health are the conditions in which people are born, grow, work, live and age, as well as the wider set of forces and systems that shape daily life conditions.
- Prematurity/LBW cause of death is a single cause of death defined as disorders related to short gestation and low birthweight (LBW), not elsewhere classified, determined by Category P07 (four codes) in the tenth International Classification of Diseases (ICD-10).
- Preterm-related cause of death is a grouping of causes of death each determined to be a direct consequence of preterm birth (44 codes from the tenth International Classification of Diseases (ICD-10)).

DATA SOURCES

- Behavioral Risk Factor Surveillance System, Centers for Disease Control and Prevention.
- IPUMS-USA, University of Minnesota, ipums.org. American Community Survey, US Census Bureau.
- National Center for Health Statistics, final natality data.
- National Center for Health Statistics, period linked birth/infant death data.

REFERENCES

- 1. Long-term health effects of preterm birth compiled by March of Dimes available at www.marchofdimes.org/complications/long-term-healtheffects-of-premature-birth.aspx
- Callaghan WM, MacDorman MF, Rasmussen SA, Qin C, Lackritz EM. The contribution of preterm birth to infant mortality rates in the United States. Pediatrics. 2006;118(4):1566-1573.
- Waitzman NJ, Jalali A, Grosse SD. Preterm birth lifetime costs in the United States in 2016: An update. Semin Perinatol. 2021 Apr;45(3):151390.
- 4. Prematurity risk factors compiled by March of Dimes available at www.marchofdimes.org/pregnancy/preterm-labor-and-birth.aspx.

